Military Use of Point of Care Ultrasound (POCUS)

Shelia C. Savell, PhD, RN[1]; Darren S. Baldwin, BS, MAA, RN[2]; Alexis Blessing, PhD[3]; Kimberly L. Medellin, BSN, RN[4]; Caroline B. Savell[5]; Joseph K. Maddry, MD[6]*

ABSTRACT

Background: Point of care ultrasound (POCUS) offers multiple capabilities in a relatively small, lightweight device to military clinicians of all types and levels in multiple environments. Its application in diagnostics, procedural guidance, and patient monitoring has not been fully explored by the Military Health System (MHS). The purpose of this narrative review of the literature was to examine the overall use of POCUS in military settings, as well as the level of ultrasound training provided. **Methods:** Studies related to use of POCUS by military clinicians with reported sensitivity/specificity, accuracy of exam, and/or clinical decision impact met inclusion criteria. After initial topical review and removal of duplicates, two authors selected 17 papers for consideration for inclusion. Four of the authors reviewed the 17 papers and determined the final inclusion of 14 studies. **Results:** We identified seven prospective studies, of which three randomized subjects to groups. Five reports described use of POCUS in patients, two used healthy volunteers, two were in simulation training environments, four used animal models to simulate specific conditions, and one used a cadaver model. Clinician subjects ranged from one to 34. Conventional medics were subjects in six studies. Four studies included special operations medics. One study included nonmedical food service inspectors. The use of ultrasound in theater by deployed consultant radiologists is described in three reports. **Conclusions:** Military clinicians demonstrated the ability to perform focused exams, including FAST exams and fracture detection with acceptable sensitivity and specificity. POCUS in the hands of trained military clinicians has the potential to improve diagnostic accuracy and ultimately care of the war fighter.

Keywords: *ultrasound; military; point of care ultrasound; POCUS*

Introduction

The first case report of hemoperitoneum diagnosis by ultrasound in a patient who experienced blunt abdominal trauma was published in 1971.[1] Subsequent studies evaluating the use of ultrasound in trauma patients led to the description of the focused assessment of sonography for trauma (FAST) exam.[2,3] The FAST exam as described in the 1993 Advanced Trauma Life Support Course became a replacement for diagnostic peritoneal lavage, which was used for initial evaluation in cases of blunt abdominal trauma.[2] The FAST assesses for intra-abdominal, intra-thoracic, and pericardial injury.[2,3] The extended FAST (eFAST) then expanded to include an evaluation for pneumothoraces, which was described by Kirkpatrick et al. in 2004.[4] The FAST/eFAST have become standard of care in trauma resuscitation and have been associated with decreased time to intervention, length of stay, cost, and complications.[5,6] The utility of the eFAST exam in the prehospital setting is described as facilitating early diagnosis and interventions, such as needle decompression for pneumothorax, as well as informing evacuation priority.[6,7] There are many other uses for POCUS, from evaluating for the presence of fracture or foreign bodies, to increasing accuracy in vascular access. POCUS can also be used in airway procedures and in focused cardiac ultrasound (FoCUS) for decision making in cardiac arrest.[8,9] Emergency ultrasound cannot replace a thorough history and exam but may quickly answer specific questions and support decision-making.[6,7]

Ultrasound technology has the potential to improve diagnostic accuracy and enhance patient care in emergency medicine and the prehospital austere setting.[6,7,11,12] Diagnostic capabilities in the combat setting may be limited, without radiography or computed tomography (CT). Portable ultrasound can be used from point of injury forward to provide support for clinical decision making by military clinicians. The Defense Advanced Research Project Administration (DARPA) awarded a grant for the development of a portable ultrasound for battlefield use in 1996. SonoSite (formally SonoSight) collaborated with the University of Washington to develop one of the first handheld ultrasound devices. In addition, several other companies began making portable ultrasound devices for civilian use.[13]

Rozanski et al.[14] describe the early use of the SonoSite 180 Plus by the US Army's 21st Combat Support Hospital (CSH) North facility in northern Iraq over a 6-month period. Multiple physician specialties performed 401 exams effectively and efficiently. The most common exam types were renal (n=174), FAST (n=69), nontrauma abdominal (n=44), and obstetric (n=40). They found the device to be versatile and reliable with clear and interpretable imaging. The use of portable ultrasound increased local diagnostic capabilities, decreased unnecessary evacuation, and was considered greatly beneficial to the far forward-deployed CSH.[14]

Military emergency medicine residency programs have incorporated extensive ultrasound training into the curriculum. In

*Correspondence to 3698 Chambers Pass, Fort Sam Houston, TX 78234 *or* Shelia.W.Savell.civ@mail.mil
[1]**Dr Shelia C. Savell**, [2]**Darren S. Baldwin**, [3]**Dr Alexis Blessing**, [4]**Kimberly L. Medellin**, [5]**Caroline B. Savell, and** [6]**Lt Col Joseph K. Maddry** are all affiliated with the US Air Force En Route Care Research Center at JBSA-Fort Sam Houston, TX. **Alexis Blessing** is also affiliated with the Oak Ridge Institute for Science and Education, Oak Ridge, TN. **Lt Col Joseph K. Maddry** is also affiliated with the Brooke Army Medical Center Department of Emergency Medicine, JBSA-Fort Sam Houston, TX.

2009, a 1-year emergency ultrasound fellowship was established and is now available at three military locations.[15] In 2011, the Joint Special Operations Medical Training Center (JSOMTC) began to incorporate a 24-hour ultrasound curriculum [Special Operator Level Clinical Ultrasound (SOLCUS) training] with didactic and hands-on training for Special Operations medics.[16] The Special Operations medical community values ultrasound as a diagnostic tool, and emphasizes the potential role of POCUS in prolonged field care.[16] The SOLCUS education project was begun to provide open-access online resources to accompany hands-on training (https://sofsono.org/).[17] The curriculum is based on the American College of Emergency Physicians (ACEP) recommendations for training of emergency physicians on emergency ultrasound. The SOLCUS program focuses on the eFAST exam and includes ocular, musculoskeletal, and soft tissue scans, as well as assessment for thrombosis.[16,17] Soon after the implementation of the SOLCUS training, the 1st Battalion, 3rd Special Forces Group (Airborne) SFG(A) deployed to Afghanistan, with nine portable ultrasound machines. Morgan et al. provides an anecdotal report of the experiences of 29 18Ds who evaluated 109 patients using POCUS.[16] The most common exams were for musculoskeletal issues (n=39), and FAST for abdominal trauma (n=34).[16]

POCUS offers multiple capabilities in a relatively small, lightweight device to military clinicians of all types and levels in multiple environments. Its application in diagnostics, procedural guidance, and patient monitoring has not been fully explored by the MHS. The purpose of this narrative review of the literature was to examine the overall use of POCUS in military settings, as well as the level of ultrasound training provided.

Methods

The authors searched PubMed, Cochrane, Scopus, Web of Science, and the Defense Technical Information Center databases using the key terms of military and ultrasound. Studies related to use of POCUS by military clinicians, with reported sensitivity/specificity, accuracy of exam, and/or clinical decision impact met inclusion criteria. After initial topical review, and removal of duplicates, two authors (SCS, AB) reviewed 40 full-text publications and selected 17 papers for inclusion consideration. Four of the authors (SCS, DSB, AB, KLM) reviewed the 17 papers and determined the final inclusion of 14 studies. Authors excluded reports that did not report sensitivity/specificity, accuracy of exam, and/or clinical decision impact, as well as literature reviews. We rated the studies using the Oxford Centre for Evidence-Based Medicine 2011 Levels of Evidence.[18] Outcomes reported in this narrative literature review include sensitivity/specificity, exam accuracy, and if the exam had impact on clinical decision making.

Results

Fourteen studies met the inclusion criteria (Table 1). However due to variance among study types, reported results, and levels of evidence, a meta-analysis was not attempted. We identified seven prospective studies, of which three used randomization of subjects to groups.[19–21,23,28–30] Five reports described use of POCUS in patients, two used healthy volunteers, and two were in simulation training environments.[19,22–25,27–32] Four studies used animal models to simulate specific conditions, and one used a cadaver model.[20,21,23,24,26] Numbers of clinician subjects ranged from 1 to 34. Conventional medics were subjects in six studies, and Special Operations medics were subjects in four studies.[19–27] One study included non-medical food service inspectors.[24] The use of ultrasound in theater by deployed consultant radiologists is described in three reports.[30–32] The levels of evidence ranged from two to four, from prospective randomized trials to a case study with four patients.[19,21,23,27]

Training

Training for five of the included studies ranged from 3 to 25 minutes and included a brief lecture, or slide show, followed by hands-on practice.[21,22,24,25,32] Five studies reported training ranging from 90 minutes to 4 hours, that also included didactic instruction followed by hands-on practice. One study reported results after Special Operations medics completed the SOLCUS course. Participants attended various components (8 to 52 hours, mean of 16.7 hours) of the SOLCUS training, prior to deployment.[27] With the exception of the three reports on radiologists, all participants had no to little prior ultrasound experience.[30–32] These studies demonstrate the ability of ultrasound naïve clinicians to successfully perform specific ultrasound exams with minimal training and practice.

Use by Military Medics and Other Nonphysicians

Conventional Army medics were the subjects of four prospective studies.[19–21,23] The medics in these studies had no prior experience or training with ultrasound. After completion of a 4-hour eFAST training, when compared to emergency medicine resident physicians, medics took longer to complete the eFAST exam (532 versus 227 seconds) but had similar diagnostic accuracy when comparing sensitivity (medics 88–95%, resident physicians 92–95%).[19] Twenty-eight medics completed a 2-hour training and identified foreign bodies in a soft tissue model with sensitivities of 73% and 78% (size dependent), and specificity of 78%.[20] A cadaver model was used to evaluate the ability of medics to detect endotracheal tube placement, after a 15-minute lecture and hands-on practice. Cadavers were randomly assigned to esophageal or tracheal tube placement for 32 participants. In an average time of 47.3 seconds, medics correctly identified tracheal placement at a rate of 72% and esophageal placement at 71%.[21] Twenty-two medics and physician assistants (number of each not specified) achieved a sensitivity of 99.2%, specificity of 95.5%, with an accuracy of 97.7% when evaluating abscesses in a tissue model.[23] Backlund et al. conducted a pilot study to assess the ability of 12 Army National Guard medics to determine cardiac activity in healthy volunteers, after a 5-minute lecture and brief hands-on training.[22] In this pilot study, 92% of the exams accurately documented the presence of cardiac activity with a mean time to completion of 5.5 seconds.[22]

Two early publications report the ability of Special Operations medics to detect fracture in a simulated model, and in four human cases.[26,27] Twenty Army Special Operations Forces (SOF) medics with minimal to no previous US experience completed a 3-minute training, and correctly identified presence or absence of long bone fracture in a simulated model (sensitivity 100%, specificity 90%).[26] US Army Special Forces medics deployed to Afghanistan evaluated 109 patients using POCUS, after completion of an average of 16.7 hours training in the SOLCUS curriculum. Medics applied their SOLCUS training

TABLE 1 *Reports of Military Use of Ultrasound*

Reference	Study Type and Aim	Subjects	Device	Training/Intervention	Results	Conclusions	LOE
Monti et al.[19]	Prospective, randomized, cohort study in the Medical Simulation Training Center at Joint Base Lewis McChord. The primary objective was to assess the impact of a 4-hour introductory training intervention on US-naive military medic participants' knowledge/ performance of the eFAST application.	34 Army medics naïve to US	Fukuda Denshi USA device with phased array (2–5 MHz) and linear array (5–12 MHz) transducer	Participants were randomized to receive either conventional, expert-led classroom didactic training or didactic training via an online, asynchronously available platform. Medic cohorts: (1) 90-min classroom didactic and 150-min hands-on scanning instruction (n=19) (2) 90-min video from SonoSim and 150-min hands-on scanning instruction (n=15)	The 50-question knowledge assessment pre/post Medic overall knowledge increased from 27% to 83%. EM residents score 92%. There was no statically significant difference between knowledge, technical performance, or diagnostic accuracy between medic groups. The medics took more than double the time to complete the exams compared to EM residents (532 vs. 227 seconds). Hemoperitoneum present sensitivity: medics 95%; EM residents 93%. Hemoperitoneum absent specificity: medics 90%; EM resident 95%. Hemopericardium present sensitivity: medics 88%; EM Residents 92%. Hemopericardium absent specificity: medics 60%; EM residents 100%.	A 4-hour introductory eFAST training intervention can effectively train conventional military medics to perform the eFAST exam. Combat medics (MOS 68W) with no previous US training compared to EM physician residents (n=20) with at least 1-month-long emergency US rotation and performed minimum of 20 eFAST exams	2
Driskell et al.[20]	Prospective, single-blinded, observational simulation to determine Army medics' accuracy performing bedside US to detect radiolucent foreign bodies in a soft-tissue hand model	28 Army medics naive to US	SonoSite M-Turbo with 13–6 MHz linear transducer	1-hour didactic and 1-hour hands-on training, used chicken model (chicken thigh in surgical glove), each medic was presented with 20 randomized models	< 2-mm FB: sensitivity 73%, specificity 78% ≥ 2-mm FB: sensitivity 78%, specificity 78%	Army medics can detect FBs in tissue models with similar sensitivities and specificities as radiologists and emergency medicine physicians in similar studies.	2
Hanlin et al.[21]	Prospective randomized trial to determine ability to detect endotracheal tube placement in a fresh human cadaver model	32 Army medics, recently completed EMT-B certification, enrolled in flight paramedic training	SonoSite M-Turbo with a 10–5 MHz linear transducer	15-min lecture on transtracheal US techniques, followed by hands-on practice	Sensitivity 66.7%, specificity 76.4% Average time 47.3 seconds; correctly identified 13/18 tracheal placements and 10/14 esophageal placements	Trainees were moderately accurate when using transtracheal US to identify ETT placement after a short educational session.	2
Backlund et al.[22]	Pilot study to assess ability of combat medics to perform a limited bedside echocardiography (BE) to determine cardiac activity in healthy volunteers	12 Army National Guard combat medics trained to level of EMT-B	Not specified	Received 5-min lecture and brief hands-on training	44 of 48 (92%) exams accurately documented presence of cardiac activity. Median time to completion–5.5 seconds	With minimal training, the majority of the medics were able to rapidly perform a focused BE on live models that was adequate to assess for the presence of cardiac activity.	3

(continues)

TABLE 1 *Cont.*

Reference	Study Type and Aim	Subjects	Device	Training/Intervention	Results	Conclusions	LOE
LaDuke et al.[23]	Single-blinded, randomized, prospective observational study to determine if nonphysician military providers could use US to detect superficial abscesses in chicken tissue models.	22 US Army medics and PAs with minimal to no prior US training	SonoSite X-Porte or SonoSite M-Turbo	Attended 30-min didactic session, followed by hands-on training	Performed 220 clinical exams and 220 US scans. US: Sensitivity 99.2%, specificity 95.5%, accuracy 97.7% Clinical exam: Sensitivity 73.5%, specificity 77.2%, accuracy 75%	With brief US training, nonphysician military medical providers can accurately detect superficial abscesses in tissue models.	2
Monti et al.[24]	Descriptive study to examine the potential for nonphysician providers to determine the absence or presence of a pneumothorax in a porcine model, with the use of a portable US machine, after receiving minimal training	22 participants (PAs, SOF, and conventional force medics, veterinary technicians, and food service inspectors) with no prior US training	SonoSite Vet (SonoSite 180 equivalent) with 10–5 MHz linear transducer	Received 10-min slide show training and orientation to US machine	Sensitivity 95.4%, specificity was 100% 21 of 22 pneumothoraces were correctly identified.	Nonphysician healthcare providers can accurately detect a pneumothorax with portable US after receiving minimal focused training.	3
Betcher et al.[25]	Proof of concept descriptive study to evaluate military trainees' ability to measure the optic nerve sheath diameter (ONSD) in healthy volunteers	23 SOCM trainees during emergency medicine rotation with minimal prior training in US	Mindray® M7	5-min lecture, followed by demonstration and 20-min practice	Compared trainee measurements to those of emergency medicine physicians: $M_{physician}$ = 0.465 mm vs. $M_{trainees}$ = 0.459 mm, p = .76	This study demonstrates that optic nerve sheath diameter measurement can be accurately performed by novice ultrasonographers after a brief training session. If validated, point-of-care optic nerve sheath diameter measurement could impact the triage of injured patients in remote areas.	3
Heiner et al.[26]	Descriptive study: simulation to evaluate the ability of 18Ds to detect the presence of long bone fracture	20 Army Special Forces (18Ds), had no or minimal prior use of US	SonoSite M-Turbo with 10–5 MHz transducer head	Received 3-min orientation and training	Sensitivity 100%, specificity 90%, 5 fracture models (turkey legs): 1 no fracture, 4 different fracture types	Using a portable US device, 18Ds were able to correctly detect the presence or absence of a simulated long bone fracture with a high degree of sensitivity and specificity.	3
Vasios et al.[27]	Case study to describe the use of portable US by 18Ds for fracture detection	29 US Army Special Forces (18Ds)–1st Battalion, 3rd Special Forces Group (Airborne) SFG(A) deployed to Afghanistan; 109 patients evaluated with US, 39 were musculoskeletal	Not specified	Received an average of 16.7 hours training using SOLCUS outline. Included FAST exam, pneumothorax detection, and musculoskeletal exam	4 cases presented – in all cases the medic correctly diagnosed fractures with US that were later confirmed by x-ray.	Highlights the potential role of the SOLCUS and the use of US by Special Operations Medics.	4

Perrier et al.[28]	Prospective study to evaluate the usefulness of point-of-care US (POCUS) performed by young military medicine residents after short practical training in the diagnosis of medical emergencies	2 military medicine residents in a French Army teaching hospital March 2015–March 2016	SonoSite M-Turbo	Received a 90-min theoretical and practical (10 USs in healthy students, 50 USs in patients with symptoms, observed by trainer) US training focused on the gallbladder, kidney, and upper urinary tract and the deep venous network of the lower extremities	Did not report sensitivity and specificity. 48 patients had US, 18 gallbladder, 16 renal, 14 lower extremity. POCUS improved diagnostic accuracy in 73% of cases, was misleading in 2% and did not contribute to 25%.	POCUS performed after clinical examination increases the diagnostic accuracy of young military medicine residents.	4
Renard et al.[29]	Prospective observational pilot study to evaluate whether the implementation of E-FAST was possible in conditions close to combat and if it changed the therapeutic and evacuation strategies	15 trainee physicians during French predeployment simulation training	Vscan (GE)	MEDICHOS (medical courses in hostile environments) internship November 2017, March and June 2018; 2-hour training on the use of the Vscan and US devices were provided, trainees to use their discretion	eFAST or POCUS exams performed on 44 of 168 (26%) simulated patients. 51% of US cases had a significant impact of therapeutic and evacuation priorities, it changed therapeutic decisions in 67% of time and evacuation priorities in 72% of time	US on the simulated battlefield was possible and useful. To confirm these results, US needs to be democratized and assessed in a real operational environment.	4
Sellon et al.[30]	Prospective questionnaire-based (6 month) study aimed to assess the usefulness of departmental diagnostic US in the remotely deployed role 2 hospital setting	Consultant radiologist at a Role 2 MTF–Op TRENTON 3	SonoSite M-Turbo with a 2.5 MHz convex probe and 10 MHz linear probe	41 departmental scans 28 July–28 December, 2017, by radiologist	In 28 of 41 (68%) cases US increased diagnostic confidence and 29% (12/41) led to a change in patient management. 1 (3%) had no clinical impact. Musculoskeletal exams had the greatest impact.	This study highlights the utility of this capability at role 2 , CT scan is not available.	4
Carter et al.[31]	Retrospective record review to determine accuracy of FAST in the deployed environment	3 consultant radiologists deployed to the Role 3 MTF, Camp Bastion, Jan–May 2014	SonoSite 4–6 MHz with curvilinear probe	Radiologists were embedded in the trauma bay	187 FAST exams performed. 169 of 187 had subsequent laparotomy or CT full body trauma scan and were included in analysis. Sensitivity 75%, Specificity 99.3% Overall accuracy 94.7% ID of intraperitoneal free fluid – PPV 96.2% and NPV 94.4%	FAST provided by the integrated radiologist as part of damage control radiology, gives the team leader rapid diagnostic information to improve decision-making and ultimately patient outcomes in the combat MTF.	3
Smith et al.[32]	Retrospective review of registry data to determine use and accuracy of FAST and CT	Attending radiologist at Role 3 MTF, Camp Bastion, July–Nov 2012	Not specified	Attending radiologists	468 casualties, 85% underwent FAST and 86.1% had CT, 34% had abdominal injury Detection of intra-abdominal injury; FAST: sensitivity 56%, specificity 98%, PPV 87%, NPV 90%, accuracy 89% CT: sensitivity 99%, specificity 99%, PPV 96%, NPV 100%, accuracy 99%	FAST and CT were useful in resuscitation care at Role 3, to enhance diagnostic sensitivity and specificity in battlefield injuries. FAST should be available in the absence of CT capability. The use of radiologists for FAST can free emergency MD to focus on other aspects of care.	

in four cases, and in all cases correctly diagnosed fractures (femur, distal fibular, phalanx, tibial), which were later confirmed by radiography.[27] In a prospective study, 23 Special Operations medic trainees underwent training to measure the optic nerve sheath diameter (ONSD) in healthy volunteers and compared their measurements to those of emergency medicine physicians.[25] After undergoing a 5-minute lecture and demonstration, medic trainees reported similar measurements in comparison to emergency medicine physicians (mean physician = 0.465 mm versus mean trainees = 0.459 mm, $p = .76$).[25] Twenty-two physician assistants, Special Operations and conventional medics, veterinary technicians, and food service inspectors (numbers of each group not specified) were able to accurately detect pneumothorax in a porcine model, after a 10-minute slide show and brief orientation to ultrasound equipment. These ultrasound naïve participants correctly identified 21 of 22 pneumothoraces, achieving a sensitivity of 95.5% and a specificity of 100%.[24]

Use by Military Physicians

Five studies reported physician use of POCUS, one simulation study, one in-garrison, and three reports of the use of ultrasound in the combat setting.[28–32] Fifteen trainee physicians completed a 2-hour ultrasound course and were provided US devices to use at will, in a simulated combat setting. The participants performed eFAST/POCUS on 44 of 168 (26%) simulated patients, and in 51% of US cases, there was a significant impact on therapeutic and evacuation priorities. Therapeutic decisions changed in 67% of cases and evacuation priorities in 72% of cases.[29] Two military medicine resident physicians performed POCUS on 48 patients in a French Army teaching hospital. POCUS improved diagnostic accuracy in 73% of cases.[28] Two studies reported the use of POCUS by radiologists, for 585 exams at a Role 3 military treatment facility (MTF) in Afghanistan.[31,32] The reported FAST sensitivity was 56 and 75%. However, specificity reached 98 and 99%, with overall accuracy of 89 and 94.4%.[31,32] A prospective 6-month survey of a consultant radiologist in a Role 2 MTF, where CT imaging capability was not available, found POCUS increased diagnostic confidence in 68% of cases and led to change in patient management in 29% of cases.[30]

Discussion

The evidence related to the use of POCUS by military clinicians is limited. This literature review included 14 published studies with mostly moderate levels of evidence. The sample sizes were small and there was not enough duplication of findings for specific exams. However, the findings consistently demonstrate the ability of military clinicians, ranging from conventional medics to physicians, to perform focused exams with moderate to high success. With minimal training, conventional medics can achieve acceptable sensitivity and specificity in FAST exams, and fracture detection. However, we found no published reports related to retention of knowledge and ability.

Conventional medics demonstrated less accuracy in the detection of foreign bodies and in confirmation of airway placement. Army Special Forces medics receive extensive training in POCUS, but only one study reported sensitivity (100%) and specificity (90%) in a simulated fracture model. Only two of five studies with physician subjects reported specificity and sensitivity. Military physicians report increased accuracy and confidence in diagnoses, as well as impact on patient management. In-theater radiologists demonstrated an overall accuracy in FAST exams of 89% to 99% (specificity 98% to 99%); however, sensitivity was only 56% to 75%.[30,31]

The findings of our review specific to military clinicians are similar to those of previous civilian literature reviews. Overall higher specificity versus sensitivity add support to the finding that POCUS has more utility in ruling in specific conditions than ruling out a condition. A systematic review of POCUS use in prehospital critical care included 27 studies and concluded POCUS is feasible and changes patient management in trauma, breathing difficulties, and in cardiac arrest. However, it is unknown if it improves patient outcomes.[7] A Cochrane review specific to use of POCUS in diagnosing thoracoabdominal injuries in patients with blunt trauma included 34 studies with 8,635 participants. The report concluded that in suspected blunt thoracoabdominal trauma, positive ultrasound findings could help guide treatment decisions.[8] Netherton et al. reviewed 75 studies (published between 1989 and 2017) on the diagnostic accuracy of the eFAST in the trauma patient.[33] This systematic review and meta-analysis suggested that eFAST was useful in ruling in pneumothorax, pericardial effusion, and intra-abdominal free fluid in trauma patients. However, the ability to "rule out" diagnoses was not supported. Similar findings were reported in a review of six papers on military use of the FAST exam.[34]

More extensive reviews of POCUS in civilian settings provide additional evidence of the potential POCUS can offer in military health care. The use of POCUS in the deployed setting has not been fully explored, however the growing evidence in support of its utility warrants ongoing study. Research needs to address which applications have the most potential to improve outcomes in combat casualty care. Training guidelines and standards should be established to determine optimal course content, length, delivery method, and the minimum number of scans for each type to achieve competence. Evidence supports image acquisition and interpretation can be taught to medical novices. However, clinical background is important for appropriate patient management.[6] Future research should consider strategies to prevent skill decay and promote knowledge retention. Research to consider the role telemedicine can play to support prehospital POCUS, the logistics of image transmission, and communication with higher-level providers is warranted. More research to determine sensitivity and specificity of various POCUS applications in the hands of military clinicians will inform best practices related to POCUS.

Further investigation into which ultrasound machine is best suited for the combat environment is required. Portable ultrasound is lightweight, easy to carry, and has minimal power requirements when compared to portable radiography, therefore increasing its utility in austere environments.[26] However, it is important to plan for anticipated conditions in the austere and resource-constrained combat theater. Considerations for device selection include: performance at high altitudes, function in extreme temperatures, protection from moisture; ultrasound gel supply, storage, and response to extreme temperatures; battery life, condition, and supply; and overall ruggedness and ability to withstand extreme conditions. SonoSite developed the first handheld device used by the military and the SonoSite M-Turbo was the most common device reported in the studies we reviewed. There are other devices with potential utility in

TABLE 2 *Portable Hand-Held Devices*

Model	Features	Size	Weight	Transducers
GE Healthcare Vscan Extend	Handheld device with dual-headed probe, store images through Wi-Fi or USB, 60-min continuous scanning on full charge, online apps available to augment studies, educational videos available, requires gel https://www.gehealthcare.com/products/ultrasound/vscan-family	**Device:** 170 × 78 × 21 mm (6.6 × 3.0 × 0.8 in) **Dual probe:** 12.9 × 3.9 × 2.8 cm (5.1 × 1.5 × 1.1 in)	**Device:** 365g (0.7 lb) **Dual probe:** 120 g (0.3 lb) **Sector probe:** 85 g **Main unit with dual probe:** 400 g	Two transducers in one probe: linear and sector
SonoSite iViz	Durable aluminum tablet with multiple transducers, cloud storage, and 64-GB flash drive, three swappable batteries each with 1-hour continuous scan time, embedded educational tools, requires gel Uses aircraft aluminum, can be dropped up to 3 feet IPX-7 rated – fully submergible in water https://www.sonosite.com/	**Tablet:** 18.3 × 11.7 × 2.7 cm (7.2 × 4.6 × 1.1 in)	**Tablet:** 570g (1.1 lb)	Curved C60v, linear 125v, linear l38v, phased p21v
Philips Lumify	Transducers attach to android devices, app-based, uses tablet as a power source, no long-term commitment, battery life depends on attached device, requires gel App based, transducers plug into devices, how-to videos Lumify System Bundle includes device https://www.usa.philips.com/healthcare/sites/lumify	**Curved transducer:** 4.5 × 11.4 cm (1.8 × 4.5 in)	**Curved transducer:** 136 g (0.3 lb)	Linear L12–4, curved c5–2, phased S4–1
Butterfly iQ	Transducer attaches to Apple mobile devices, built-in battery, wireless charging, unlimited cloud storage, uses silicon chip, does not use Piezo crystal technology, 2 hours of continuous scanning on full charge, no gel required Anodized aluminum body, thermally efficient, educational videos available https://www.butterflynetwork.com/	**Transducer:** 185 – 56 – 35 mm (7.2 × 2.2 × 1.4 in)	**Transducer:** 313 g (0.7 lb)	Single transducer emulates any kind of transducer
Clarius C3 Convex	App based, wireless, does not require Internet access to operate Handheld device with three probes in one, works on iOS and Android educational videos available Has magnesium shell, waterproof, and withstands drops up to 1 m; 60-min battery power, three swappable batteries 2–6 MHz, max depth 32 cm https://clarius.com/	**Device:** 167 × 99 × 42 mm (6.6 × 3.9 × 1.6 in)	**Device:** 540 g (1.2 lb)	Three clip-on tips to scan entire body

Note: Some data from Canepa and Harris (2019).[2]

combat care (Table 2). The end-users should be consulted to determine desired device characteristics. Forty combat medics compared a novel US finger transducer to a conventional transducer in a simulation study.[35] Diagnostic accuracy was similar between transducers; however, the mean completion times were better with the conventional transducer (304 versus 358 seconds; $p = .03$). In addition, the medics scored the conventional transducer higher for ease of use.[35] Additional research comparing available technology needs to be conducted.

There are limitations to the conclusions that can be drawn from the findings of a narrative literature review. The method for determining inclusion/exclusion may be somewhat subjective and lead to selection bias. The focus of this literature review on military personnel only, yielded studies with small sample sizes and variations in outcome measures. Studies of civilians with similar roles and training may have larger sample sizes and outcomes that are more congruent. However, it is important to describe the available evidence for this unique population, in order to lay the foundation for future study.

Well-designed prospective, randomized studies with larger sample sizes, as well as longitudinal designs, would yield more meaningful data to determine level of training required. These can also inform sustainment training. There is a need to conduct prospective and retrospective studies, designed to evaluate the use of POCUS by military clinicians. These should include evaluation of sensitivity and specificity, as well as patient outcomes.

Conclusions

Military clinicians demonstrated the ability to perform focused exams, including FAST exams and fracture detection, with acceptable sensitivity and specificity. POCUS in the hands of trained military clinicians has the potential to improve diagnostic accuracy and ultimately care of the war fighter.

Presentation

This literature review was presented in online poster format at the San Antonio Universities and Military Research Forum (SURF), June 2020. It was also presented in poster format at the 2020 Special Operations Medical Association Scientific Assembly.

Funding

Defense Health Agency J9. This research was supported in part by an appointment to the Postgraduate Research Participation Program at the US Army Institute of Surgical Research administered by the Oak Ridge Institute for Science and

Education through an interagency agreement between the US Department of Energy and USAISR.

Disclosures

The views expressed are solely those of the authors and do not reflect the official policy or position of the US Army, US Navy, US Air Force, the Department of Defense, or the US Government. The authors have no financial relationship relevant to this article to disclose.

Author Contributions

JKM and SCS conceived the study concept. JKM obtained funding. SCS, CBS and DSB completed the literature search coordinated and collected the data, and AB reviewed and summarized findings in table. SCS wrote the first draft, and all authors read, edited, and approved the final manuscript.

References

1. **Kristensen JK, Bue B, Kühl E.** Ultrasonic scanning in the diagnosis of splenic haematomas. *Acta.Chir Scand*. 1971;137(7):653–657.
2. **Rippey JCR, Royse AF**. Ultrasound in trauma. *Best Pract Res Clin Anaesthesiol*. 2009;23(3):343–362.
3. **Rozycki GS, Ochsner MG, Jaffin JH, Champion HR**. Prospective evaluation of surgeons' use of ultrasound in the evaluation of trauma patients. *J Trauma*. 1993:34(4):516–526.
4. **Kirkpatrick AW, Sirois M, Laupland KB, et al.** Hand-held thoracic sonography for detecting post-traumatic pneumothoraces: the Extended Focused Assessment with Sonography for Trauma (eFAST). *J Trauma*. 2004;57(2):288–295.
5. **Bloom BA, Gibbons RC.** Focused Assessment with Sonography for Trauma. [Updated 2020 Jul 31]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan–. Available from: https://www.ncbi.nlm.nih.gov/books/NBK470479/
6. **Canepa CA, Harris NS.** Ultrasound in austere environments. *High Alt Med Bio*. 2019; 20(2):103–111.
7. **Botker MT, Jacobson L, Rudolph SS, Knudsen L.** The role of point of care ultrasound in prehospital critical care: a systematic review. *Scand J Trauma Resus Emerg Med*. 2018;26:51.
8. **Gharahbaghian L, Anderson KL, Lobo V, et al.** D. Point-of-care ultrasound in austere environments: a complete review of its utilization, pitfalls, and technique for common applications in austere settings. *Emerg Med Clin N Am*. 2017;35:409–441.
9. **Nations JA, Browning RF.** Battlefield applications for handheld ultrasound. *Ultrasound Q*. 2011;27:171–176.
10. **Stengel D, Leisterer J, Ekkernkamp S, et al.** Point-of-care ultrasonography for diagnosing thoracoabdominal injuries in patients with blunt trauma. *Cochrane Database Syst Rev.* 2018;12(12): CD012669.
11. **Ketelaars R, Reijnders G, van Geffenm GJ, et al.** ABCDE of prehospital ultrasonography: a narrative review. *Crit Ultrasound J*. 2018;10(17).
12. **Nelson BP, Melnick ER, Li J.** Portable ultrasound for remote environments, part II: current indications. *J Emerg Med*. 2011;40 (3):313–21.
13. **Nelson BP, Melnick ER, Li J.** Portable ultrasound for remote environments, part I: Feasibility of field deployment. *J Emerg Med*. 2011;40(2):190–197.
14. **Rozanski TA, Edmondson JM, Jones SB.** Ultrasonography in a forward-deployed military hospital. *Mil Med*. 2005;170(2):99–102.
15. **Hile DC, Morgan AR, Laselle BT, Bothwell JD.** Is point-of-care ultrasound accurate and useful in the hands of military medical technicians? A review of the literature. *Mil Med*. 2012;177(8):983–987.
16. **Morgan AR, Vasios WN, Hubler DA, Benson PJ.** Special operator level clinical ultrasound: an experience in application and training. *J Spec Oper Med*. 2010;10(2):16–21.
17. **Hampton K, Vasios WN, Loos PE.** SOLCOS: update on point-of-care ultrasound in Special Operations medicine. *J Spec Oper Med*. 2016;16(1):58–61.
18. **Howick J, Chalmers I, Glasziou P, et al.** The 2011 Oxford CEBM Evidence Levels of Evidence (Introductory Document). Oxford Centre for Evidence-Based Medicine. https://www.cebm.ox.ac.uk/resources/levels-of-evidence/levels-of-evidence-introductory-document. Accessed 5 March 2021.
19. **Monti JD, Perreault MS.** Impact of a 4-hour introductory eFAST training intervention among ultrasound-naïve U.S. military medics. *Mil Med*. 2020;185(5-6):e601-e608.
20. **Driskell DL, Gillum JB, Monti JD, Cronin A.** Ultrasound evaluation of soft-tissue foreign bodies by US army medics. *J Med Ultrasound*. 2018;26:147–152.
21. **Hanlin ER, Zelenak J, Barakat J, Anderson KL.** Airway ultrasound for the confirmation of endotracheal tube placement in cadavers by military flight medic trainees – A pilot study. *Am J Emerg Med*. 2018;36(9):1711–1714.
22. **Backlund BH, Bonnett CJ, Faragher JP, Haukoos JS, et al.** Pilot study to determine the feasibility of training Army National Guard medics to perform focused cardiac ultrasonography. *Prehosp Emerg Care*. 2010;14(1):118–123.
23. **LaDuke M, Monti J, Cronin A, Gillum B.** Ultrasound detection of soft tissue abscesses performed by non-physician U.S. Army medical providers naïve to diagnostic sonography. *Mil Med*. 2017;182(3):e1825-e1830.
24. **Monti JD, Younggren B, Blankenship R.** Ultrasound detection of pneumothorax with minimally trained sonographers: a preliminary study. *J Spec Oper Med*. 2009;9(1):43–46.
25. **Betcher J, Becker TK, Stoyanoff P.** Military trainees can accurately measure optic nerve sheath diameter after a brief training session. *Mil Med Res*. 2018;5(42).
26. **Heiner JD, Baker BL, McArthur TJ.** The ultrasound detection of simulated long bone fractures by US Army Special Forces medics. *J Spec Oper Med*. 2010;10(2):7–9.
27. **Vasios WN, Hubler DA, Lopez RA, Morgan AR.** Fracture detection in a combat theater: four cases comparing ultrasound to conventional radiology. *J Spec Oper Med*. 2010;10(2):11–15.
28. **Perrier P, Leyral J, Thabouillot O, et al.** Usefulness of point-of-care ultrasound in military medical emergencies performed by young military medicine residents. *J R Army Med Corps*. 2019;0:1–4.
29. **Renard A, Martinet C, Cungi PJ, et al.** Is E-FAST possible and useful on the battlefield? A feasibility study during medical courses in hostile environment (MEDICHOS): preliminary results. *J R Army Med Corps*. 2019;165:338–341.
30. **Sellon E, Durdle S, Bailey D.** Assessing the utility of ultrasound in the role 2 hospital setting. *J R Army Med Corps*. 2019;0:1–4.
31. **Carter NJ, Gay D.** FAST in the deployed military setting. *J R Army Med Corps*. 2018;164:332–334.
32. **Smith IM, Naumann DN, Marsden MER, et al.** Scanning and war: utility of FAST and CT in the assessment of battlefield abdominal trauma. *Ann Surg*. 2015;262(2):389–296.
33. **Netherton S, Milenkovic V, Taylor M, Davis PJ.** Diagnostic accuracy of eFAST in the trauma patient: a systematic review and meta-analysis. *CJEM*. 2019;21(6):727–738.
34. **Qi X, Jing Tian J, Sun R, et al.** Focused assessment with sonography in trauma for assessment of injury in military settings: a meta-analysis. *Balkan Med J*. 2020;37:3–8.
35. **Ahern BJ, Monti JD, Naylor JF, et al.** U.S. Army combat medic eFAST performance with a novel versus conventional transducers: a randomized, crossover trial. *Mil Med*. 2020;185(Suppl 1): 19–24.

The Special Operations Medical Association's Official Journal

JSOM

JOURNAL *of* SPECIAL OPERATIONS MEDICINE™

Summer 2021
Volume 21, Edition 2

THE JOURNAL FOR OPERATIONAL MEDICINE AND TACTICAL CASUALTY CARE

Inside this Issue:

- FEATURE ARTICLES: Tourniquet Practice Models
- Atherosclerosis in Elite Special Operations Forces
- 23.4% Hypertonic Saline for TBI
- The Effect of Airdrop on Fresh Whole Blood
- Military POCUS › Unconventionally Acquired Brain Injury
- Prehospital Trauma Registry After-Action Reviews in Afghanistan
- Telemedicine Capabilities of Special Operations
- Targeted Intervention in Patients With mTBI
- Back Pain in Italian Helicopter Aircrews › TXA Use in TEMS Providers
- CASE REPORTS: TXA Use in TEMS Providers › Infectious Myositis › Bacteria on Female Urinary Diversion Devices
- IN BRIEF: Far-Forward Blood Donation
- ONGOING SERIES: Human Performance Optimization, Infectious Disease, Injury Prevention, Law Enforcement & Tactical Medicine, Research: How To, Veterinary Medicine, TCCC Updates, *and more!*

Dedicated to the Indomitable Spirit, Lessons Learned & Sacrifices of the SOF Medic

A Peer Reviewed Journal That Brings Together The Global Interests of Special Operations' First Responders